

Learning Mixtures of Separated non-Spherical Gaussians

Sanjeev Arora*
Dept of Computer Science
Princeton University.

Ravi Kannan†
Dept of Computer Science
Yale University

Abstract

Mixtures of gaussian (or normal) distributions arise in a variety of application areas. Many heuristics have been proposed for the task of finding the component gaussians given samples from the mixture, such as the *EM algorithm*, a local-search heuristic from Dempster, Laird and Rubin (1977). These do not provably run in polynomial time.

We present the first algorithm that provably learns the component gaussians in time that is polynomial in the dimension. The Gaussians may have arbitrary shape; but they must satisfy a “separation condition” which places a lower bound on the distance between the centers of any two component Gaussians. The mathematical results at the heart of our proof are “distance concentration” results —proved using isoperimetric inequalities— which establish bounds on the probability distribution of the distance between a pair of points generated according to the mixture.

We also formalize the more general problem of max-likelihood fit of a gaussian mixture to unstructured data.

1 Introduction

Finite mixture models are ubiquitous in a host of areas that use statistical techniques, including AI, computer vision, medical imaging, psychology, geology etc. (see [15, 23]). A mixture of distributions $\mathcal{D}_1, \mathcal{D}_2, \dots$ with mixing weights w_1, w_2, w_3, \dots (where $\sum_i w_i = 1$) is the distribution in which a sample is produced by first picking a component distribution —the i th one is

*Supported by NSF Grants CCR-0098180 and CCR-0205594, Alfred Sloan Fellowship, and a David and Lucille Packard Fellowship. arora@cs.princeton.edu

†Supported in part by NSF Grant CCR-9820850. kannan@cs.yale.edu

picked with probability w_i — and then producing a sample from that distribution. In many applications the component distributions are multivariate gaussians.

Given samples from the mixture distribution, how can one learn (i.e., reconstruct) the component distributions and their mixing weights? The most popular method is probably the EM algorithm of Dempster, Laird and Rubin (1977). EM is a local search heuristic that tries to converge to a *maximum likelihood* description of the data by trying to cluster sample points according to which gaussian they came from. Though moderately successful in practice, it often fails to converge or gets stuck in local optima. Much research has gone into fixing these problems, but has not yet resulted in an algorithm that provably runs in polynomial time. A second known technique is called “projection pursuit” in Statistics (Huber [12]). In this, one projects the samples into a random low dimensional space and then in the projected space, try to do the clustering (exhaustively) exploiting the low dimensionality.

We note that some combinatorial problems seemingly related to learning gaussian mixtures are NP-hard. For instance, Megiddo [18] shows that it is NP-hard to decide, given a set of points in \Re^n , whether the points can be covered by two unit spheres. This problem seems related to learning a mixture of two spherical gaussians.

Nevertheless, one may hope that when the data is generated from the mixture of gaussians (as opposed to being unstructured as in Megiddo’s result) then the algorithm could use this structure in the data. Recently, Dasgupta (1999) [5] took an important step in this direction by showing how a mixture of k identical gaussians could be learned in polynomial time provided the gaussians are “spherelike” (their probability mass is concentrated in a thin spherical shell) and their centers are “well-separated.” (Such separation conditions correspond to a non-degeneracy assumption: if the mixture contains two identical gaussians whose centers are arbitrarily close, then they cannot be distinguished even in principle.)

Though Dasgupta’s algorithm is a good first step, it leaves open the question whether one can design algorithms that require weaker assumptions on the gaussians. This is the topic of the current paper: our algorithms make no assumption about the shape of the gaussians but they require the gaussians to be “well-separated.” Even for the special case of spherical gaussians, our result improves Dasgupta’s (and a result of Dasgupta and Schulman [6] that is independent of our work). We describe our results in more detail in Section 2.3, and compare them to other work.

We also define a more general problem of gaussian fitting, whereby we

make no assumptions about the data and have to fit the mixture of k gaussians that maximizes the log-likelihood it assigns to the dataset (see Section 2.1). We use techniques developed in context of approximation algorithms to design algorithms for one of the problems (see Section 4). The exact problem is NP-hard.

2 Definitions and Overview

The univariate distribution $N(\mu, \sigma)$ on \mathfrak{R} has the density function $f(x) = (\sqrt{2\pi}\sigma)^{-1} \exp(-\frac{(x-\mu)^2}{2\sigma^2})$. It satisfies $E[(x - \mu)^2] = \sigma^2$. The analogous distribution in \mathfrak{R}^n is the *axis-aligned* gaussian $N(\bar{\mu}, \bar{\sigma})$ where $\bar{\mu}, \bar{\sigma} \in \mathfrak{R}^n$ and the density function is the product distribution of $N(\mu_1, \sigma_1), N(\mu_2, \sigma_2), \dots, N(\mu_n, \sigma_n)$. A random sample (x_1, x_2, \dots, x_n) satisfies $E[\sum_i (x_i - \mu_i)^2] = \sum_i \sigma_i^2$. (Similarly, $E[\sum_i (x_i - \mu_i)^2 / \sigma_i^2] = n$.)

A general gaussian in \mathfrak{R}^n is obtained from an axis-aligned gaussian by applying an arbitrary rotation. Specifically, its probability density function has the form

$$F_{Q,p}(x) = \frac{1}{(2\pi)^{n/2} \prod_i \sqrt{\lambda_i(Q)}} \exp(-(x-p)^T Q^{-1}(x-p)/2). \quad (1)$$

where Q is an $n \times n$ positive definite matrix with eigenvalues $\lambda_1(Q), \dots, \lambda_n(Q) > 0$, and $p \in \mathfrak{R}^n$ is the *center*. Since Q can be rewritten as $R^{-1} \text{diag}(\lambda_i(Q)) R$ where R is a rotation, the quantities $\lambda_i(Q)$ play the same role as the variances σ_i^2 in the axis-aligned case. From our earlier discussion, $E[(x-p)^T(x-p)]$ is $\sum_i \lambda_i(Q)$ and $E[(x-p)^T Q^{-1}(x-p)] = \int_x F_{Q,p}(x) (x-p)^T Q^{-1}(x-p) = n$.

For any finite sample of points in \mathfrak{R}^n we can try to fit a gaussian by computing their *variance-covariance matrix*. Let x_1, x_2, \dots, x_N be N points in \mathfrak{R}^n in general position (i.e., we assume that they do not all lie on a hyper-plane). Let X be the $n \times N$ matrix whose columns are the vectors $x_1 - q, x_2 - q, \dots, x_N - q$, where $q = \frac{1}{N}(x_1 + x_2 + \dots + x_N)$ is the sample mean. Then the variance-covariance matrix $A = \frac{1}{N} X X^T$; note that it is positive definite by definition.

This fit may, of course, be poor for an arbitrary point set. But if the points are sampled from gaussian $F_{G,p}$, then $F_{A,q}$ converges rapidly to $F_{G,p}$: with $\Omega(n \log n / c_\epsilon)$ samples (where c_ϵ is a constant depending upon ϵ), $F_{A,q}$ is with high probability a $(1 + \epsilon)$ -fit to $F_{G,p}$ in every direction [3, 22] in the sense that $|q - p|^2 \leq \epsilon \sum_i \lambda_i(Q)$ and $|Gv|(1 - \epsilon) \leq |Av| \leq (1 + \epsilon)|Gv|$ for every unit length vector v . (The proof is fairly elementary for $\Omega(n^2/\epsilon^2)$ samples.)

Spherical and spherelike gaussians In an axis-aligned gaussian with center at the origin and with variances $\sigma_1^2, \dots, \sigma_n^2$, the quantity $\sum_i x_i^2 / \sigma_i^2$ is the sum of n iid variables from $N(0, 1)$ so this sum is tightly concentrated about its mean n . In a *spherical* gaussian, all σ_i 's are the same, so even $\sum_i x_i^2$ is tightly concentrated. (These observations go back to Borel.) More generally, $E[\sum_i x_i^2] = \sum_i \sigma_i^2$. If the σ_i 's are not "too different", then distance-concentration results (similar to Lemma 5 below) show that that almost all of the probability mass is concentrated in a thin spherical shell of radius about $(\sum_i \sigma_i^2)^{1/2}$; such gaussians may be thought of as *spherelike*. Roughly speaking, if $\text{radius} / \sigma_{\max} = \Omega(\log n)$, then the gaussian is spherelike. Known algorithms (such as Dasgupta [5]) work only for such spherelike gaussians. By contrast, here, we wish to allow Gaussians of all shapes.

2.1 Max likelihood learning

Now we formalize the learning problems. Consider a mixture of gaussians $(w_1, F_1, w_2, F_2, \dots, w_m, F_m)$ in \mathfrak{R}^n , where the w_i 's are the mixing weights. With any point $x \in \mathfrak{R}^n$, one can associate m numbers $(F_i(x))_{i=1, \dots, m}$ corresponding to the probabilities assigned to it by the various gaussians according to formula (1). For any sample $S \subseteq \mathfrak{R}^n$ this imposes a natural partition into m blocks: each point $x \in S$ is labeled with an integer $l(x) \in \{1, \dots, m\}$ indicating the distribution that assigns the highest probability to x . (Ties are broken arbitrarily.) The *likelihood* of the sample is

$$\prod_{x \in S} F_{l(x)}(x).$$

It is customary to work with the logarithm of this likelihood, called the *log-likelihood*.

Thus one may mean several things when talking about "learning mixtures of gaussians" [21]. The following is the most general notion.

Definition 1 (Max likelihood fit) *In the Max Likelihood fit problem, we are given an arbitrary sample $S \subseteq \mathfrak{R}^n$, a number k , and desire the gaussian mixture with k components that maximizes the likelihood of S .*

2.2 Classification Problem

Now we define the sub-case of the learning problem when the data is assumed to arise from an unknown mixture of k gaussians, where k is known.

Definition 2 (Classification problem) *In the classification problem, we are given an integer k , a real number $\delta > 0$ and a sample S generated from an unknown mixture F_1, F_2, \dots, F_k of k gaussians in \mathbb{R}^n , where the mixing weights w_1, w_2, \dots, w_k are also unknown. The goal is to find the “correct” labeling for each point in S , (upto permutation), namely to partition S into k subsets such that with probability at least $1 - \delta$, the partition of S is exactly into the subsets of samples drawn according to each F_i .*

Viewing the unknown mixture as a “source” we may view this as the “source learning” problem. Note that once we know the partition, we can immediately obtain estimates of the unknown gaussians and their mixing weights.

So, the Classification problem has a stronger hypothesis than the maximum-likelihood problem in that it assumes that the data came from a mixture. It also then requires the result to satisfy the stronger requirement that it is exactly the partition into the actual S_1, S_2, \dots, S_k , where, S_i was generated according to the Gaussian F_i . (We abuse notation here slightly; we can only know the real S_i upto a permutation of their indices. But to avoid extra notation, we will say the partition has to be S_1, S_2, \dots, S_k .)

2.3 Our results

Our main result concerns the classification problem. Clearly, the problem has no unique solution if the gaussians in the mixture are allowed to be arbitrarily close to each other. We will assume a certain separation between the centers of the Gaussians. The required separation is an important consideration and will be motivated in detail in section 3.2. Here we will just state it and mention two important features of it.

Notation : First, we introduce some notation which we will use throughout. We let p_1, p_2, \dots, p_k denote the (unknown) centers respectively of the k Gaussians F_1, F_2, \dots, F_k comprising the mixture; the maximum variance of F_i in any direction will be denoted $\sigma_{i,max}$. We will denote by R_i the “median radius” of F_i ; R_i has the property that the F_i -measure of a ball of radius R_i around p_i is exactly $1/2$. Henceforth, we will reserve the word “radius” of a Gaussian to mean its median-radius. Here is our formal definition of separation.

Definition 3 *For any $t > 0$, we say that the mixture is t -separated if*

$$\begin{aligned} |p_i - p_j|^2 \geq & -|R_i^2 - R_j^2| + 500t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) \\ & + 100t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2) \quad \forall i, j \end{aligned} \quad (2)$$

We point out here quickly two features of this definition. Firstly, if two Gaussians F_i, F_j are both spherical of the same radii ($R_i = R_j$), then the required separation is $O^*(R_i/n^{1/4})$. Secondly, if F_i, F_j are still spherical, but, if $R_j > R_i$, the term $-|R_i^2 - R_j^2|$ is negative and makes the separation required lesser. Indeed if $R_j = \Omega^*((1 + (1/\sqrt{n}))R_i)$, then the two Gaussians F_i, F_j are allowed to be concentric!¹

Theorem 1 *There is a polynomial-time algorithm for the classification problem. The algorithm needs to know a lower bound w_{min} on the mixing weights, and the number s of sample points required is $O(n^2 k^2 \log(kn^2)/(\delta^2 w_{min}^6))$. The gaussians may have arbitrary shape but have to be t -separated where $t = O(\frac{\log s}{\delta})$.*

We also present an approximation algorithm for a special case of the max-likelihood fit problem.

Theorem 2 *There is a polynomial-time approximation algorithm for the max likelihood fit problem in \mathbb{R}^n when the gaussians to be fitted to the data have to be spherical of equal radii (the radius and the centers of the k gaussians have to be determined by the algorithm). There is a fixed constant c such that the algorithm produces a solution whose log-likelihood is at least the best possible minus c .*

The algorithm of Theorem 2 is combinatorial and appears in Section 4. We note even this sub-case of the maximum likelihood fit problem is at least as hard as the clustering problem k -median (sum-of-squares version with steiner nodes), which is NP-hard [9]. Indeed, our algorithm is obtained by reducing to the k -median algorithm of [4] (recent more efficient k -median algorithms would also work). We feel that this way of viewing the learning problem as an approximation problem may be fruitful in other contexts.

2.4 Comparison with other results

The algorithm in Dasgupta [5] makes the following assumptions: (i) all the Gaussians have the same variance-covariance matrix Σ ; (ii) the maximum and minimum eigenvalues - σ_{max}^2 and σ_{min}^2 respectively of Σ satisfy $\frac{\sigma_{max}}{\sigma_{min}} \in O(\sqrt{n}/\log k)$ and (iii) the centers of any 2 of the k Gaussians are at distance (at least) $\Omega(\sqrt{n}\sigma_{max})$ apart.

Dasgupta and Schulman [6] showed that the EM algorithm learns (and indeed does so in just two rounds) a mixture of **spherical** gaussians F_1, F_2, \dots, F_k ,

¹The superscript * over O, Ω indicates that we have omitted logarithmic in n factors.

where F_i has radius R_i (the R_i may be different.) They require now only a separation between centers of F_i, F_j of $\Omega((R_i + R_j)/n^{1/4})$. (This amount of separation ensures among other things that the densities are “non-overlapping” - i.e., there are k disjoint balls, each containing the samples picked according to one F_i .)

As mentioned, our result is stronger in two ways. First, we allow Gaussians of arbitrary (and different) variance-covariance matrices and secondly we allow densities to overlap, or even be concentric. More, specifically, the term $-|R_i^2 - R_j^2|$ (which is non-positive) can make the minimum required separation negative (and so a vacuous requirement) in some cases; it allows the centers to be close (or even coincide) if the radii are very different. This allows a “large feature” to have a identifiable smaller “feature” buried inside. For the case dealt with by [6], their requirement is the same as ours (since in this case, $R_i \approx \sqrt{n}\sigma_{i,max}$) but for this term and logarithmic factors and thus their result essentially follows as a special case of ours. For the case dealt with by [5], our requirement is again weaker than that paper’s but for logarithmic factors (since $\sqrt{n}\sigma_{i,max} \in O(R_i)$). After the first appearance of our paper [2], Vempala and Wong [25] have improved the separation requirement to essentially the optimal one for the special case of spherical Gaussians — $|p_i - p_j| = \Omega((R_i + R_j)/\sqrt{n})$. Their spectral technique is entirely different from ours.

3 Algorithm for Classification problem

First we fix notation for the rest of the section. We are given a set S of samples, picked according to an unknown mixture $w_1 F_1 + w_2 F_2 + \dots w_k F_k$ of Gaussians $F_1, F_2, \dots F_k$. The known quantities are k , and a number w_{min} that is a lower bound on the w_i ’s. We have to decompose S as $S = S_1 \cup S_2 \cup \dots \cup S_k$, where S_i are the samples that were generated using F_i .

Section 3.1 describes the algorithm at an intuitive level. This description highlights the need for a “well-separation” condition on the gaussians, which we explain in Section 3.2. The description also highlights the need for “distance concentration” results for gaussians, which are then proved in Section 3.3. In Section 3.5 we formally describe the algorithm and prove its correctness.

3.1 Algorithm overview

The algorithm uses distance-based clustering, meaning that we repeatedly identify some sample point x and some distance r , and all sample points

in $B(x, r)$ all put into the same cluster. Such distance-based clustering is not new and it appears in many heuristics including [5, 6]. The choice of x, r is of course the crucial new element we provide. Since distance-based methods seem restrictive at first glance, the surprising part here is that we get provable results which subsume previously known provable results for any algorithm. This power arises from a “bootstrapping” idea, whereby we learn a little about the gaussians from a coarse examination of the data and then bootstrap from that information to find a better clustering.

In general, distance-based clustering is most difficult when the gaussians have different shapes and sizes, and overlap with each other (all of which we allow). It is easy to see why: a sample point from gaussian F_i might be closer to some sample points of another gaussian F_j than to all the sample points of F_i . One crucial insight in our algorithm is that this is unlikely to happen if we look at the Gaussian with the smallest radius in the mixture; hence we should use clustering to first identify this gaussian, and then iterate to find the remaining gaussians.

Now we give an overview of the algorithm. Let F_1 be the gaussian of smallest radius. Using our distance-concentration results, we can argue that for any $x \in S_1$, there is a r such that (i) $B(x, r) \cap S = S_1$. (ii) There is a “sizable” gap after r , namely, the annulus $B(x, r') \setminus B(x, r)$ for some r' noticeably larger than r contains no samples from any S_j for $j > 1$. (iii) There is no spurious large gaps before r , which would confuse the algorithm.

Even after proving the above statements, the design of the algorithm is still unclear. The problem is to figure out the size of the gap between where S_1 ends and $S \setminus S_1$ begins, so we know when to stop. (Note: there will be gaps before r ; the point is that they will be smaller than the one after r .) Our separation condition ensures that the gap between S_1 and the other S_j is $\Omega(\sigma_{1,max})$; so we need an estimate of $\sigma_{1,max}$. We get such an estimate by bootstrapping. We show that: (i) if we have any fraction f of the samples in S_1 , then we may estimate $\sigma_{1,max}$ to a factor of $O(1/f^2)$ with high probability. We use this to get a rough estimate β of $\sigma_{1,max}$. Using β , we increment the radius in steps which are guaranteed to be less than $\sigma_{1,max}$ (which ensures that we do not step over the “gap” into another S_j) until we observe a gap; by then, we have provably picked up *most* of S_1 . Now we use this to better estimate $\sigma_{1,max}$ and then incrementing the radius by another $\Omega(\sigma_{1,max})$, we capture all of S_1 . (The guaranteed gap ensures that we do not get any points from any other Gaussian while we increment the radius.)

To make all the above ideas rigorous, we need appropriate distance concentration results which assert that the distance between certain pairs of sample points considered as a random variable is concentrated around a

certain value. Some distance concentration results— at least for spherical or spherelike gaussians— were known prior to our work, showing a sharp concentration around the mean/median. However, for the current algorithm we also need concentration around values that are quite far from the mean/median. For example, to show the nonexistence of “spurious gaps,” we have to show that if a ball of radius r centered at a sample point $x \in S_i$ has F_i -measure, say, exactly $1/4$, then for a small $\delta > 0$, the ball of radius $r + \delta$ with x as center has F_i -measure at least 0.26 . If such a result failed to hold, then we might see a “gap” (an annulus with no sample points) and falsely conclude that we have seen all of S_i . Such concentration results (around values other than the median/mean) are not in general provable by the traditional moment-generating function approach. We introduce a new approach in this context: Isoperimetric inequalities (see Theorem 3). Our method does not always prove the tightest possible concentration results, but is more general. For example, one may derive weaker concentration results for general log-concave densities via this method (see [17]).

3.2 Separation condition and its motivation

Now we motivate our separation condition, which is motivated by the exigencies of distance-based clustering. Consider the very special case of spherical Gaussians F_i, F_j with $R_i = R_j$. Suppose x, x', y are independent samples, x, x' picked according to F_i and y according to F_j . Lemma 5 will argue that with high probability (we will use \approx here to mean that the two sides differ by at most $O(R_i^2/\sqrt{n})$) :

$$|x - p_i|^2 \approx R_i^2$$

and similar concentration results for $|x' - p_i|, |y - p_j|$. It is an intuitive fact that $x - p_i, x' - p_i, p_i - p_j, y - p_j$ will all be pairwise nearly orthogonal (a sample from a spherical gaussian is almost orthogonal to any fixed direction with high probability). So, one can show that

$$|x - x'|^2 \approx |x - p_i|^2 + |p_i - x'|^2 \approx 2R_i^2 \quad (3)$$

$$|x - y|^2 \approx |x - p_i|^2 + |p_i - p_j|^2 + |y - p_j|^2 \approx 2R_i^2 + |p_i - p_j|^2. \quad (4)$$

(The first assertion is proved rigorously in greater generality in Lemma 7 and the second one in Lemma 8.) Thus, it is clear that if $|p_i - p_j|^2$ is at least $\Omega(R_i^2/\sqrt{n})$, then with high probability $|x - x'|$ and $|x - y|$ will lie in different ranges. (Aside: One can also show a sort of converse with different constants, since the concentration results we get are qualitatively tight. However, we

will not establish this, since it is not needed.) This inter-center separation then is

$$O(R_i/n^{1/4}).$$

Our separation condition for this case is indeed this quantity, upto a $\log n$ factor. A weaker separation condition would be to require a separation of $\Omega(R_i/\sqrt{n})$; at this separation, one can still show that with high probability the hyperplane orthogonal to the line joining the centers at its mid-point has all the samples of one Gaussian on one side and the samples of the other Gaussian on the other side. An algorithm to learn under this condition would be a stronger result than our distance-based algorithm in this case. After the conference version of our paper appeared, Vempala and Wang [25] have indeed developed a learning algorithm under this weaker separation for the case of spherical Gaussians using spectral techniques.

3.3 Concentration results using isoperimetric inequalities

Suppose we have some probability density F in \mathbb{R}^n and a point x in space. For proving distance concentration results, we would like to measure the rate of growth / decline of $F(B(x, r))$ as a function of r . This will be provided by the Isoperimetric inequality (see Corollary 4).

Theorem 3 [14] *Let $F(x) = e^{-x^T A^{-1} x} g(x)$ be a function defined on \mathbb{R}^n , where A is a positive definite matrix whose largest eigenvalue is σ_{max}^2 and $g(x)$ is any positive real valued log-concave function. Suppose ν is a positive real and we have a partition of \mathbb{R}^n into three sets K_1, K_2, K_3 so that for all $x \in K_1, y \in K_2$, we have $|x - y| \geq \nu$. Then*

$$\int_{K_3} F(x) dx \geq \frac{2\nu e^{-\nu^2}}{\sqrt{\pi}} \frac{1}{\sigma_{max}} \min \left(\int_{K_1} F(x) dx, \int_{K_2} F(x) dx \right).$$

The phrase ‘‘isoperimetric inequality’’ has come to mean a lower bound on the surface area of a set in terms of its volume. If K_1 is fixed and we define K_3 to be the set of points not in K_1 which are at distance at most ν from some point in K_1 and define $K_2 = \mathbb{R}^n \setminus (K_1 \cup K_3)$, then as ν goes to zero, K_3 tends to the boundary surface of K_1 and the above theorem can be shown to yield a lower bound on the surface integral of F over this surface. We will make this connection rigorous below for the context we need. Such isoperimetric inequalities for general log-concave measures over multi-dimensional sets were first proved for use in establishing rapid convergence to the steady state of certain Markov chains for approximating volumes of

convex sets and for sampling according to log-concave measures. [8], [16], [1]. The proof of Theorem 3 uses a specialization of the above techniques to the case of Gaussians, where we get better results.

Corollary 4 *We borrow notation from Theorem 3 and also assume that $F(\mathfrak{R}^n) = 1$.*

(i) *If a ball $B(x, r)$ has $F(B(x, r)) \leq 1/2$, then*

$$\frac{d(\ln(F(B(x, r))))}{dr} \geq \frac{2}{\sqrt{\pi}\sigma_{max}}.$$

(ii) *If a ball $B(x, r)$ has $F(B(x, r)) \geq 1/2$, then*

$$\frac{d(\ln(1 - F(B(x, r))))}{dr} \leq \frac{-2}{\sqrt{\pi}\sigma_{max}}.$$

Remark: The corollary says that $\ln(F(B(x, r)))$ grows at a rate of $\Omega(1/\sigma_{max})$ until $F(B(x, r))$ is $1/2$, and then $\ln(1 - F(B(x, r)))$ declines at a rate of at a rate of $\Omega(1/\sigma_{max})$. Intuitively, it is easy to see that this would lead to distance concentration results since once we increase (decrease) r by $O(\sigma_{max})$ from its median value, the mass outside $B(x, r)$ (inside $B(x, r)$) is small. The first lemma below (Lemma 5) is derived exactly on these lines; the subsequent three Lemmas 6,7,8 discuss the distances between different samples from the same and from different Gaussians.

Proof. Let ν be an infinitesimal. Then,

$$\begin{aligned} \frac{d(\ln(F(B(x, r))))}{dr} &= \frac{1}{F(B(x, r))} \frac{d(F(B(x, r)))}{dr} \\ &= \lim_{\nu \rightarrow 0} \frac{1}{\nu F(B(x, r))} [F(B(x, r + \nu)) - F(B(x, r))]. \end{aligned}$$

Now we let $K_1 = B(x, r)$ and $K_2 = \mathfrak{R}^n \setminus B(x, r + \nu)$ and apply the Theorem above to get the first assertion of the corollary. The second assertion follows similarly. \square

Lemma 5 *Suppose F is a general Gaussian in \mathfrak{R}^n with maximum variance in any direction σ , radius R and center p . Then for any $t > 0$, we have*

$$F(\{x : R - t\sigma \leq |x - p| \leq R + t\sigma\}) \geq 1 - e^{-t}.$$

Proof. For any $\gamma > 0$, let $F(B(p, \gamma)) = g(\gamma)$. Then for $\gamma < R$, we have by Corollary (4), that

$$\frac{dg(\gamma)}{d\gamma} \geq \frac{g(\gamma)}{\sigma}.$$

So, $\frac{d \ln(g(\gamma))}{d\gamma} \geq \frac{1}{\sigma}$. Integrating from γ to R , we get that

$$F(B(p, \gamma)) \leq \frac{1}{2} e^{-(R-\gamma)/\sigma}.$$

For $\gamma > R$, isoperimetry implies that

$$\frac{dg(\gamma)}{d\gamma} \geq \frac{1-g(\gamma)}{\sigma} \implies \frac{d(\ln(1-g(\gamma)))}{d\gamma} \leq \frac{-1}{\sigma}.$$

Again integrating from R to γ , we get $1-g(\gamma) \leq (1/2)e^{-(\gamma-R)/\sigma}$. Combining the two, the lemma follows. \square

Lemma 6 *Let F, p, R, σ be as in Lemma 5 and suppose z is any point in space. Let $t \geq 1$. If x is picked according to F , we have that with probability at least $1 - 2e^{-t}$,*

$$\begin{aligned} (R + t\sigma)^2 + |z - p|^2 + 2\sqrt{2}\sqrt{t}|z - p|\sigma &\geq |x - z|^2 \\ &\geq ((R - t\sigma)^+)^2 + |z - p|^2 - 2\sqrt{2}\sqrt{t}|z - p|\sigma \end{aligned} \quad (5)$$

where $(R - t\sigma)^+$ is $R - t\sigma$ if this quantity is positive and 0 otherwise.

Proof. We have

$$\begin{aligned} |x - z|^2 &= ((x - p) + (p - z)) \cdot ((x - p) + (p - z)) \\ &= |x - p|^2 + |p - z|^2 + 2(x - p) \cdot (p - z). \end{aligned} \quad (6)$$

Now $2(x - p) \cdot (p - z)$ is a normal random variable with mean 0 and variance at most $4|p - z|^2\sigma^2$, so the probability that $|2(x - p) \cdot (p - z)|$ is greater than $2\sqrt{2}\sqrt{t}|z - p|\sigma$ is at most e^{-t} . From lemma (5), we have that $R - t\sigma \leq |x - p| \leq R + t\sigma$ with probability at least $1 - e^{-t}$. Combining these two facts, the current lemma follows. \square

Lemma 7 *Suppose F, p, R, σ as in Lemma 5. Suppose x, y are independent samples each picked according to F . Then for any $t \geq 1$, with probability at least $1 - 3e^{-t}$, we have*

$$2R^2 - 8t\sigma R \leq |x - y|^2 \leq 2(R + 2t\sigma)^2.$$

Proof. We may assume that x is picked first and then y (independently). Then from lemma (5), with probability $1 - e^{-t}$, we have $R - t\sigma \leq |x - p| \leq R + t\sigma$. From Lemma 6, (once x is already picked), with probability at least $1 - 2e^{-t}$, we have $(R + t\sigma)^2 + |x - p|4\sqrt{t}\sigma + |x - p|^2 \geq |x - y|^2 \geq R^2 - 2Rt\sigma - 4|x - p|\sqrt{t}\sigma + |x - p|^2$. Both the conclusions hold with probability at least $1 - 3e^{-t}$, whence we get :

$$|x - y|^2 \leq (R + t\sigma)^2 + 4t\sigma(R + t\sigma) + (R + t\sigma)^2 \leq 2(R + 2t\sigma)^2.$$

For the lower bound on $|x - y|^2$, first note that if $R \leq 4t\sigma$, then $2R^2 - 8t\sigma \leq 0$, so the lower bound is obviously valid. So we may assume that $R > 4t\sigma$. Thus, $|x - p| \geq 3t\sigma$ and under this constraint, $|x - p|^2 - 4\sqrt{t}\sigma|x - p|$ is an increasing function of $|x - p|$. So, we get

$$|x - y|^2 \geq R^2 - 2Rt\sigma - 4t\sigma(R - t\sigma) + (R - t\sigma)^2,$$

which yields the lower bound claimed. \square

Lemma 8 *Let $t \geq 1$. If x is a random sample picked according to F_i and y is picked independently according to F_j , with F_i, F_j satisfying the separation condition (2), then with probability at least $1 - 6e^{-t}$, we have*

$$\begin{aligned} |x - y|^2 &\geq 2 \min(R_i^2, R_j^2) + 60t(\sigma_{i,max} + \sigma_{j,max})(R_i + R_j) \\ &\quad + 30t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2) \end{aligned} \quad (7)$$

Proof. Assume without loss of generality that $R_i \leq R_j$. Applying Lemma 6, we get that with probability at least $1 - 2e^{-t}$, we have :

$$|y - p_i|^2 \geq R_j^2 - 2t\sigma_{j,max}R_j + |p_i - p_j|^2 - 2\sqrt{2}\sqrt{t}|p_i - p_j|\sigma_{j,max}. \quad (8)$$

Claim 1

$$|y - p_i|^2 \geq R_i^2 + 154t(\sigma_{i,max} + \sigma_{j,max})(R_i + R_j) + 30t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2). \quad (9)$$

Proof. Case 1 : $R_j^2 \geq R_i^2 + 250t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) + 30t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2)$. Note that $|p_i - p_j|^2 - 4\sqrt{t}\sigma_{j,max}|p_i - p_j| + 4t\sigma_{j,max}^2 \geq 0$. So, $|p_i - p_j|^2 - 4\sqrt{t}\sigma_{j,max}|p_i - p_j| \geq -4t\sigma_{j,max}^2$. Plugging this into (8), and using the case assumption, we get

$$\begin{aligned} |y - p_i|^2 &\geq R_i^2 + 250t(R_i + R_j)(\sigma_{j,max} + \sigma_{i,max}) \\ &\quad + 30t^2(\sigma_{j,max}^2 + \sigma_{i,max}^2) - 2t\sigma_{j,max}R_j - 4t\sigma_{j,max}^2. \end{aligned}$$

It is easy to see that $R_j \geq (2/3)\sigma_{j,max}$ - this is because R_j^2 is clearly at least the median value of $(u \cdot x)^2$ under F_j , where u is the direction achieving $\sigma_{j,max}$; now it is easy to see that for the 1-dimensional Gaussian - $u \cdot x$, the median value of $(u \cdot x)^2$ is at least $2/3$ times the variance by direct calculation. Plugging in $\sigma_{j,max}^2 \leq (3/2)R_j\sigma_{j,max}$ into the above inequality, we easily get the claim.

Case 2 : $R_j^2 < R_i^2 + 250t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) + 30t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2)$. Then by the separation condition, we have

$$|p_i - p_j|^2 \geq 250t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) + 70t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2)$$

Now, since $|p_i - p_j|^2 - 2\sqrt{2}\sqrt{t}\sigma_{j,max}|p_i - p_j|$ is an increasing function of $|p_i - p_j|$ for $|p_i - p_j| \geq 2\sqrt{2}\sqrt{t}\sigma_{j,max}$, we have

$$\begin{aligned} |p_i - p_j|^2 - 2\sqrt{2}\sqrt{t}\sigma_{j,max}|p_i - p_j| &\geq 250t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) \\ &\quad - 2\sqrt{2}\sqrt{t}\sigma_{j,max}(16\sqrt{t}\sqrt{R_j + R_i}\sqrt{\sigma_{j,max} + \sigma_{i,max}}) \\ &\quad + 70t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2) + 9t(\sigma_{j,max} + \sigma_{i,max}) \\ &\geq 156t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) \\ &\quad + 34t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2) \quad (\text{using } \sigma_{j,max} \leq (3/2)R_j). \end{aligned}$$

Putting this into (8), we get :

$$|y - p_i|^2 \geq R_i^2 - 2t\sigma_{j,max}R_j + 156t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) + 34t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2),$$

which yields the claim in this case. \square

Imagine now y already having been picked and x being picked independently of y . Applying Lemma 6, we get that with probability at least $1 - 2e^{-t}$, we have

$$\begin{aligned} |x - y|^2 &\geq R_i^2 - 2R_it\sigma_{i,max} + |y - p_i|^2 - 2\sqrt{2}\sqrt{t}\sigma_{i,max}|y - p_i| \\ &\geq R_i^2 - 2R_it\sigma_{i,max} + R_i^2 + 154t(\sigma_{i,max} + \sigma_{j,max})(R_i + R_j) \\ &\quad + 30t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2) - 2\sqrt{2}\sqrt{t}\sigma_{i,max}(R_i + 13\sqrt{t}\sqrt{\sigma_{i,max} + \sigma_{j,max}}\sqrt{R_i + R_j}) \\ &\quad + \sqrt{30}t(\sigma_{i,max} + \sigma_{j,max}) \end{aligned}$$

(because under condition (9), $|y - p_i|^2 - 2\sqrt{2}\sqrt{t}\sigma_{i,max}|y - p_i|$ is an increasing function of $|y - p_i|$)

$$\begin{aligned} &\geq 2R_i^2 + (154 - 2 - 2\sqrt{2}(1 + 13\sqrt{3/2} + (1.5)\sqrt{30}))t(\sigma_{i,max} + \sigma_{j,max})(R_i + R_j) \\ &\quad + 30t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2), \end{aligned}$$

from which the lemma follows. \square

3.4 Warm-up: case of Spherical Gaussians

As a consequence of our concentration results we first present our algorithm for the simple case when all the F_i are spherical. In this case, $\sigma_{i,max} \approx R_i/\sqrt{n}$, where the error is small enough that our calculations below are valid. Choosing $t = \Omega(\log(|S|/\delta))$ as before, it is easy to see that from the distance concentration results that with high probability ,

$$|x - y|^2 \in \left[2R_i^2 \left(1 - \frac{4t}{\sqrt{n}}\right) \quad , \quad 2R_i^2 \left(1 + \frac{5t}{\sqrt{n}}\right) \right] \quad \forall x, y \in S_i \quad \forall i \quad (10)$$

and by appropriately choosing the constant in the definition of t -separation we can also ensure that with high probability there is a positive constant $c' > 12$ such that

$$|x - y|^2 \geq 2 \min(R_i^2, R_j^2) + \frac{c't(R_i + R_j)^2}{\sqrt{n}} \quad (11)$$

$$\forall x \in S_i \quad \forall y \in S_j, \quad \forall i \neq j.$$

For each pair, $x, y \in S$, find $|x - y|$ and suppose x_0, y_0 is a pair (there may be several) at the minimum distance. Then from (10) and (11), it follows that if $x_0 \in S_i$, then for all $y \in S_i$, $|x - y| \leq (1 + \frac{3t}{\sqrt{n}})|x_0 - y_0|$ and furthermore for all $z \in S \setminus S_i$, $|x - z| \geq (1 + \frac{6t}{\sqrt{n}})|x_0 - y_0|$. So, we may identify S_i by $S \cap B(x_0, |x_0 - y_0|(1 + \frac{3t}{\sqrt{n}}))$. Having thus found S_i , we may peel it off from S and repeat the argument. The important thing here is that we can estimate the radius of the ball —namely, $|x_0 - y_0|(1 + \frac{3t}{\sqrt{n}})$ —from observed quantities; this will not be so easily the case for general Gaussians.

3.5 The general case

Now we consider the case when the gaussians may not be spherical or even spherelike. Let $\delta > 0$ be the probability of failure allowed. We are given a set of samples S drawn according to an unknown mixture of Gaussians $w_1 F_1 + w_2 F_2 + \dots w_k F_k$; but we are given a $w_{min} > 0$ with $w_i \geq w_{min}$ for all i . We assume that $|S| \geq 10^7 n^2 k^2 \log(kn^2)/(\delta^2 w_{min}^6)$. In what follows, we choose

$$t = \frac{100 \log |S|}{\delta}.$$

The Algorithm

Initialization : $T \leftarrow S$.

1. Let $\alpha > 0$ be the smallest value such that a ball $B(x, \alpha)$ of radius α centered at some point in $x \in T$ has at least $3w_{\min}|S|/4$ points from T . [This will identify a Gaussian F_i with approximately the least radius.]
2. Find the maximum variance of the set $Q = B(x, \alpha) \cap T$ in any direction. I.e., find

$$\beta = \max_{w:|w|=1} \frac{1}{|Q|} \sum_{y \in Q} (w \cdot y - w \cdot (\frac{1}{|Q|} \sum_{z \in Q} z))^2.$$

[This β is our first estimate of σ_{max} . Note that computing β is an eigenvalue computation, and an approximate eigenvalue suffices.]

3. Let $\nu = \sqrt{\frac{w_{\min}\beta}{8}}$. [We will later show that $\nu \leq \sigma_{max}$; so increasing the radius in steps of ν ensures that we do not miss the “gap” between the S_i that x belongs to and the others.] Find the least positive integer s such that [we will later prove that such a s exists].

$$B(x, \alpha + s\nu) \cap T = B(x, \alpha + (s - 1)\nu) \cap T.$$

4. Let $Q' = B(x, \alpha + s\nu) \cap T$. As in Step 3, find the maximum variance β' of the set Q' in any direction. [We will prove that this β' gives a better estimate of σ_{max} .]
5. Remove $B(x, \alpha + s\nu + 3\sqrt{\beta'}(\log |S| - \log \delta + 1)) \cap T$ from T . [We will show that the set so removed is precisely one of the S_i .]
6. Repeat with until T is empty.

Remark 1 *Ball $B(x, \alpha + s\nu)$ will be shown to contain all but $w_{\min}/(10w_i)$ of the mass of the Gaussian F_i we are dealing with; the bigger radius of $B(x, \alpha + s\nu + 3\sqrt{\beta'}(\log |S| - \log \delta + 1))$ will be shown to include all but $\delta/(10|S|^2)$ of the mass of F_i . This will follow using Isoperimetry. Then we may argue that with high probability all of S_i is now inside this ball. An easier argument shows that none of the other S_j intersect this ball.*

Now we prove why this works as claimed. Let $\delta > 0$ be the probability of failure allowed. We choose

$$t = \frac{100 \log |S|}{\delta}.$$

We will now show using the distance concentration results that several desirable events — described below in (12),(13),(14), (15), (16), (17) and

(18) happen, each with probability at least $1 - \frac{\delta}{10}$. **We will assume from now on that the conditions (12) through (18) hold after allowing for the failure probability of at most $7\delta/10$.** The bottom line is that the sample is very likely to represent the mixture accurately: the component gaussians are represented essentially in proportion to their mixing weights; the number of samples in every sphere and half-space is about right; and so forth.

First, since $|S_i|$ can be viewed as the sum of $|S|$ Bernoulli independent 0-1 random variables, where each is 1 with probability w_i , we have (using standard results - eg. Hoeffding's inequality which asserts that for s i.i.d. Bernoulli random variables X_1, X_2, \dots, X_s with $\text{Prob}(X_i = 1) = q$, for all real numbers $\alpha > 0$, $\text{Prob}(|\sum_{i=1}^s X_i - sq| \geq \alpha) \leq 2e^{-\alpha^2 q/4s}$) that with probability at least $1 - \delta/10$,

$$1.1w_i|S| \geq |S_i| \geq (0.9)w_i|S| \forall i. \quad (12)$$

For each $i, 1 \leq i \leq k$ and each $x \in S_i$, let $\eta(x)$ be the least positive real number such that

$$F_i(B(x, \eta(x))) \geq 1 - \frac{\delta}{10|S|^2}.$$

Now, we assert that with probability at least $1 - \frac{\delta}{10}$,

$$\forall i, 1 \leq i \leq k, \forall x \in S_i, \quad S_i \subseteq B(x, \eta(x)). \quad (13)$$

To see this, focus attention first on one particular $x \in S$, say $x \in S_i$. We may imagine picking x as the first sample in S and then independently picking the others. Since then x is fixed, $\eta(x)$ and also $B(x, \eta(x))$ are fixed; so from the lower bound on $F_i(B(x, \eta(x)))$, it follows that $\text{Prob}(S_i \subseteq B(x, \eta(x))) \geq 1 - \delta/(10|S|)$. From this, we get (13).

We have from Lemma 7 that with probability at least $1 - \delta/10$, the following is true for each $i, 1 \leq i \leq k, \forall x, y \in S_i$:

$$2R_i^2 - 8t\sigma_{i,max}R_i \leq |x - y|^2 \leq 2(R_i + 2t\sigma_{i,max})^2. \quad (14)$$

Further, from Lemma 8, we have that with probability at least $1 - \frac{\delta}{10}$,

$$\begin{aligned} \forall i, j \leq k, i \neq j, \forall x \in S_i, \forall y \in S_j, \\ |x - y|^2 \geq 2 \min(R_i^2, R_j^2) + 60t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) \\ + 30t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2). \end{aligned} \quad (15)$$

Next, we wish to assert that certain spherical annuli centered at sample points have roughly the right number of points. Namely,

$$\begin{aligned} & \forall i, \quad \forall x, y, z \in S_i \text{ letting } A = B(x, |x - y|) \setminus B(x, |x - z|), \\ & \text{we have} \quad \left| \frac{|S_i \cap A|}{|S_i|} - F_i(A) \right| \leq \frac{w_{min}^{5/2}}{160}. \end{aligned} \quad (16)$$

We will only sketch the routine argument for this. First, for a particular triple x, y, z in some S_i , we may assume that these points x, y, z were picked first and the other points of the sample are then being picked independently. So for the other points, the annulus is a fixed region in space. Then we may view the rest as Bernoulli trials and apply Hoeffding's inequality. The above follows from the fact that the Hoeffding upper bound multiplied by $|S|^3$ (the number of triples) is at most $\delta/10$.

Next, we wish to assert that for every half-space in space, we have about the correct number of points. For this, we note that the VC (Vapnik-Chervonenkis) dimension [24] of half-spaces is n and from their theorem, it follows that with probability at least $1 - \delta/10$, we have

$$\begin{aligned} & \forall i \quad \forall \text{ half spaces } H \\ & ||S_i \cap H| - |S_i|F_i(H)| \leq w_{min}|S_i|/100 \end{aligned} \quad (17)$$

From Lemma 12, (to come), it follows that with probability at least $1 - \delta/10$, we have

$$\begin{aligned} & \forall \text{ unit length vectors } w, \quad \forall i, \\ & \frac{1}{|S_i|} \sum_{x \in S_i} (w \cdot (x - p_i))^2 \leq 2\sigma_{i,max}^2. \end{aligned} \quad (18)$$

Lemma 9 *Each execution of steps 1-5 removes precisely one of the S_i*

The lemma will be proved by induction on the number of executions of the loop. Suppose we have finished $l - 1$ executions and are starting on the l th .

Let P be the set of j such that S_j has not yet been removed. (By the inductive assumption at the start of the loop, T is the union of $S_j, j \in P$.)

Lemma 10 *Suppose $x \in S$ is the center of the ball $B(x, \alpha)$ found in the l th execution of step (1) of the algorithm and suppose x belongs to S_i (i*

unknown to us). Then,

$$B(x, \alpha) \cap S \subseteq S_i \quad (19)$$

$$\begin{aligned} |x - y|^2 &\geq 2R_i^2 + 50t(\sigma_{i,max} + \sigma_{j,max})(R_i + R_j) \\ &\quad + 20t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2) \quad \forall y \in S_j \forall j \neq i, j \in P. \end{aligned} \quad (20)$$

Proof. For any $j \in P$, and all $y, z \in S_j$, we have from (14) that $|z - y|^2 \leq 2(R_j + 2t\sigma_{j,max})^2$. Thus, a ball of radius $\sqrt{2}(R_j + 2\sigma_{i,max})$ with y as center would qualify in step (1) of the algorithm by (12). So, by definition of α in that step, we must have

$$\alpha \leq \sqrt{2}(R_j + 2t\sigma_{j,max}) \forall j \in P. \quad (21)$$

If now $B(x, \alpha)$ contains a point z from some S_j , $j \neq i$, by the inductive assumption in lemma (9), we must have $j \in P$. Then by (15), we have

$$\alpha^2 \geq 2 \min(R_i^2, R_j^2) + 60t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) + 30t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2),$$

which contradicts (21) (noting that (21) must hold for both i, j). This proves (19).

Now, from the lower bound of (14), it follows that

$$\alpha^2 \geq 2R_i^2 - 8R_i\sigma_{i,max}t.$$

So, from (21), it follows that

$$2R_j^2 \geq 2R_i^2 - 8t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) - 8t^2\sigma_{j,max}^2 \forall j \in P.$$

Thus from (15), we get that for $y \in S_j$; $j \neq i$,

$$\begin{aligned} |x - y|^2 &\geq 2R_i^2 - 8t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) - 8t^2\sigma_{j,max}^2 \\ &\quad + 60t(R_i + R_j)(\sigma_{i,max} + \sigma_{j,max}) + 30t^2(\sigma_{i,max}^2 + \sigma_{j,max}^2), \end{aligned} \quad (22)$$

from which (20) follows. \square

Now we can show that β is a rough approximation to $\sigma_{i,max}$.

Claim 2 *The β, Q computed in step 2 of the algorithm satisfy*

$$\frac{2|S_i|}{|Q|}\sigma_{i,max}^2 \geq \beta \geq \frac{|Q|^2}{4|S_i|^2}\sigma_{i,max}^2.$$

Proof. For any unit length vector w , we have by (18) :

$$\sum_{x \in Q} (w \cdot (x - p_i))^2 \leq \sum_{x \in S_i} (w \cdot (x - p_i))^2 \leq 2|S_i|\sigma_{i,max}^2$$

Since this holds for every w , and the second moment about the mean is less than or equal to the second moment about p_i , we have that $\beta \leq 2\frac{|S_i|}{|Q|}\sigma_{i,max}^2$. This proves the upper bound on β .

Let u be the direction of the maximum variance of F_i . We wish to assert that the variance of Q along u is at least $|Q|\sigma_{i,max}/|S_i|$. To this end, first note that for any reals γ_1, γ_2 , with $\gamma_1 > 0$, we have :

$$\begin{aligned} \text{Prob}_{F_i}(\gamma_2 - \gamma_1 \leq x \cdot u \leq \gamma_2 + \gamma_1) &= \frac{1}{2\sqrt{\pi}\sigma_{i,max}} \int_{\gamma_2 - \gamma_1}^{\gamma_2 + \gamma_1} e^{-r^2/2\sigma_{i,max}^2} dr \\ &\leq \gamma_1/(\sqrt{\pi}\sigma_{i,max}). \end{aligned}$$

Let $\gamma_2 = \frac{1}{|Q|} \sum_{x \in Q} (u \cdot x)$ and let $\gamma_1 = \frac{|Q|}{|S_i|}\sigma_{i,max}$. Then the strip $H = \{x : \gamma_2 - \gamma_1 \leq u \cdot x \leq \gamma_2 + \gamma_1\}$ satisfies $F_i(H) \leq \gamma_1/(\sqrt{\pi}\sigma_{i,max})$. So by (17)

$$|S_i \cap H| \leq |S_i|F_i(H) + \frac{w_{min}|S_i|}{100} \leq 3|Q|/4 \text{ using } |Q| \geq \frac{3}{4}w_{min}|S|.$$

So, we have that

$$\begin{aligned} \frac{1}{|Q|} \sum_{x \in Q} (u \cdot x - \gamma_2)^2 &\geq \frac{1}{|Q|} \frac{|Q|}{4} \frac{|Q|^2}{|S_i|^2} \sigma_{i,max}^2 \\ &= \frac{1}{4} \sigma_{i,max}^2 \frac{|Q|^2}{|S_i|^2}, \end{aligned}$$

from which the lower bound on β obviously follows. \square

Corollary 11 *The β computed in step 2) of the algorithm satisfies :*

$$\frac{4}{w_{min}} \sigma_{i,max}^2 \geq \beta \geq \frac{1}{8} w_{min}^2 \sigma_{i,max}^2.$$

Proof. Since $|Q| \geq 3w_{min}|S|/4$, the Claim above implies the corollary. \square

Now since we are increasing the radius in steps of ν which is at most $\sigma_{i,max}$ (Corollary 11), and Lemma 10 implies that samples from other gaussians are ‘‘far’’ from x compared to the samples of F_i , (this is immediate for the first iteration; for subsequent iterations of the algorithm, we use the

already established inductive step of Lemma 9 which says that what is left is precisely the union of some S_i and then apply Lemma 10) it follows that there is an annulus around x which contains no samples. Thus we have proved the existence of the desired s in step 3), and also that

$$B(x, \alpha + s\nu) \cap S \subseteq S_i.$$

The trouble of course is that such a gap may exist even inside of S_i , so $B(x, \alpha + s\nu)$ may not contain all of S_i . To complete the induction we have to argue that the steps (4) and (5) will succeed in identifying *every* point of S_i . For $\gamma > 0$, let

$$g(\gamma) = F_i(B(x, \gamma)).$$

From $B(x, \alpha + s\nu) \cap T = B(x, \alpha + (s-1)\nu) \cap T$, (see step 3 of the algorithm), we get using (16) that

$$g(\alpha + s\nu) - g(\alpha + (s-1)\nu) \leq \frac{w_{min}^{5/2}}{160}.$$

So, there exists a $\gamma' \in [(\alpha + (s-1)\nu), (\alpha + s\nu)]$ with

$$\left(\frac{dg(\gamma)}{d\gamma} \right)_{\gamma=\gamma'} \leq \frac{w_{min}^{5/2}}{160\nu} \leq \frac{w_{min}}{20\sigma_{i,max}}.$$

(If not, integration would contradict the previous inequality.) Thus Isoperimetry (Corollary 4) implies that

$$g(\alpha + s\nu) \geq 1 - \frac{w_{min}}{10} \quad \text{OR} \quad g(\alpha + (s-1)\nu) \leq 0.1w_{min}.$$

The latter is impossible since even the α -radius ball contains at least $3|S|w_{min}/4$ points. This implies that $g(\alpha + s\nu) \geq 0.9$ and now again using (16), we see that $|Q'| \geq 0.8|S_i|$ (note that Q' is found in step 4). Thus from Claim 2, (noting that the proof of works for any subset of S_i), we get that β' is a fairly good approximation to $\sigma_{i,max}$:

$$2.5\sigma_{i,max}^2 \geq \beta' \geq 0.16\sigma_{i,max}^2. \quad (23)$$

From the definition of s in step (3) of the algorithm, it follows that there is some $y \in S_i$ with $|x - y| \geq \alpha + (s-2)\nu$. So, from (14), we have $\alpha + (s-2)\nu \leq \sqrt{2}(R_i + 2t\sigma_{i,max})$. So, we have

$$\begin{aligned} \alpha + s\nu + 3\sqrt{\beta'} \left(\log \frac{|S|}{\delta} + 1 \right) &\leq \sqrt{2}(R_i + 2t\sigma_{i,max}) + 2\sigma_{i,max} \\ &+ 4\sigma_{i,max} \left(\log \frac{|S|}{\delta} + 1 \right). \end{aligned} \quad (24)$$

Thus from (20), no point of S_j , $j \in P \setminus \{i\}$ is contained in $B(x, \alpha + s\nu + 3\sqrt{\beta'}(\log \frac{|S|}{\delta} + 1))$. So the set removed from T in step 5) is a subset of S_i .

Finally, using $g(\alpha + s\nu) \geq 9/10$, and Isoperimetry (Corollary 4), we see that

$$g(\alpha + s\nu + 3\sqrt{\beta'}(\log \frac{|S|}{\delta} + 1)) \geq 1 - \frac{\delta}{10|S|^2},$$

which implies that $\eta(x) \leq \alpha + s\nu + 3\sqrt{\beta'}(\log \frac{|S|}{\delta} + 1)$. Thus, by (13), all of S_i is in $B(x, \alpha + s\nu + 3\sqrt{\beta'}(\log \frac{|S|}{\delta} + 1))$. This completes the inductive proof of correctness. \square

Now we prove a lemma that was used above when we estimated σ_{\max} .

Lemma 12 *Suppose F is a (general) Gaussian in \mathfrak{R}^n . If L is a set of independent identically distributed samples, each distributed according to F , then with probability at least $1 - \frac{\delta}{10}$, we have (with $\epsilon = 20n(\sqrt{\log n} + \sqrt{\log(1/\delta)})/\sqrt{|L|}$), every vector w satisfies*

$$\begin{aligned} E_F((w \cdot (x - E_F(x)))^2)(1 - \epsilon) &\leq E_S((w \cdot (x - E_F(x)))^2) \\ &\leq E_F((w \cdot (x - E_F(x)))^2)(1 + \epsilon), \end{aligned} \quad (25)$$

where E_S denotes the “sample mean”, i.e., it stands for $\frac{1}{|L|} \sum_{x \in L}$.

Proof. We may translate by $-E_F(x)$ and without loss of generality assume that $E_F(x)$ is the origin. Suppose Q is the square root of the inverse of the variance-covariance matrix of F . We wish to prove for all vectors w :

$$E_F((w \cdot x)^2)(1 - \epsilon) \leq E_S((w \cdot x)^2) \leq E_F((w \cdot x)^2)(1 + \epsilon).$$

Putting $Q^{-1}w = u$ (noting that Q is nonsingular and symmetric), this is equivalent to saying for all vectors u ,

$$E_F((u \cdot (Qx))^2)(1 - \epsilon) \leq E_S((u \cdot (Qx))^2) \leq E_F((u \cdot (Qx))^2)(1 + \epsilon).$$

But Qx is a random sample drawn according to the standard normal, so it suffices to prove the statement for the standard normal. To prove it for the standard normal, we proceed as follows. First for each coordinate i , we have that $E_F(|x_i|^2) = 1$ and using properties of the standard one-dimensional normal density, for any real $s > 0$,

$$\text{Prob}(|E_S(|x_i|^2) - 1| \leq s) \geq 1 - e^{-|L|s^2/4}.$$

Now consider a pair $i, j \in \{1, 2, \dots, n\}$, where $i \neq j$. The random variable $x_i x_j$ has mean 0 and variance 1. $E_S(x_i x_j)$ being the average of N i.i.d. samples (each not bounded, but we may use the properties of the normal density again) concentrated about its mean :

$$\text{Prob}(E_S|x_i x_j| \leq s) \geq 1 - e^{-|L|s^2/100}.$$

Putting $s = 10 \frac{\sqrt{\log n}}{\sqrt{|L|}}$, we see that all these $O(n^2)$ upper bounds hold simultaneously with probability at least $1 - \delta/n^8$.

Thus we have that the ‘‘moment’’ of inertia matrix M of S whose i, j entry is $E_S(x_i x_j)$ has entries between $1 - \frac{1}{2}\epsilon$ and $1 + \frac{1}{2}\epsilon$ on its diagonal and the sum of the absolute values of the entries in each row is at most $\epsilon/2$. Thus by standard Linear Algebra (basically arguments based on the largest absolute value entry of any eigenvector), we have that the eigenvalues of M are between $1 - \epsilon$ and $1 + \epsilon$ proving what we want. \square

4 Max-likelihood estimation

Now we describe an algorithm for max-likelihood fit of a mixture of k spherical gaussians of equal radius to (possibly) unstructured data. First we derive a combinatorial characterization of the optimum solution in terms of the k -median (*sum of squares, steiner version*) problem. In this problem, we are given M points $x_1, x_2, \dots, x_M \in \mathfrak{R}^n$ in \mathfrak{R}^n and an integer k . The goal is to identify k points p_1, p_2, \dots, p_k that minimize the function

$$\sum_{i=1}^M |x - p_{c(j)}|^2, \tag{26}$$

where $p_{c(j)}$ is the point among p_1, \dots, p_k that is closest to j and $|\cdot|$ denotes Euclidean distance.

Theorem 13 *The mixture of k spherical gaussians that minimizes the log-likelihood of the sample is exactly the solution to the above version of k -median.*

Proof. Recall the density function of a spherical gaussian of variance σ (and radius $\sigma\sqrt{n}$) is

$$\frac{1}{(2\pi\sigma)^{n/2}} \exp(-|x - p|^2 / 2\sigma^2).$$

Let $x_1, x_2, \dots, x_M \in \mathfrak{R}^n$ be the points. Let p_1, p_2, \dots, p_k denote the centers of the gaussians in the max-likelihood solution. For each data point x_j let $p_{c(j)}$ denote the closest center. Then the mixing weights of the optimum mixture w_1, w_2, \dots, w_k are determined by considering, for each i , the fraction of points whose closest center is p_i .

The log-likelihood expression is obtained by adding terms for the individual points to obtain the following.

$$- \left[\text{Constant} + \frac{Mn}{2} \log \sigma + \sum_j \frac{|x_j - p_{c(j)}|^2}{2\sigma^2} \right].$$

The optimum value $\hat{\sigma}$ is obtained by differentiation :

$$\hat{\sigma}^2 = \frac{2}{Mn} \sum_j |x_j - p_{c(j)}|^2, \quad (27)$$

which simplifies the log-likelihood expression to

$$\text{Constant} + \frac{Mn}{2} \log \hat{\sigma} + \frac{Mn}{4}.$$

Thus the goal is to minimize $\hat{\sigma}$, which from (27) involves minimizing the familiar objective function from the sum-of-square version of the k -median problem. \square

We indicate how to use known results about k -median to provide a constant *additive factor* approximation to the log likelihood. Charikar et al. [4] provide an $O(1)$ approximation to the k -median problem with sum-of-square distance. They do not indicate if their algorithm works when the centers are not one of the sample points. However, triangle inequality implies that picking the center to be one of the sample points changes the objective of the k -median problem by at most a factor 4. Hence we obtain a constant factor approximation to $\hat{\sigma}^2$, and hence an approximation to $\log(\hat{\sigma})$ that is correct within an $O(1)$ *additive factor*. More efficient algorithms for k -median are now known, so there is some hope that the observations of this section may lead to some practical learning algorithms.

5 Conclusions

Several open problems remain. The first concerns solving the classification problem for gaussians with significant overlap. For example, consider mixtures of spherical gaussians with pairwise inter-center distance only

$O(\max\{\sigma_1, \sigma_2\})$. In this case, a constant fraction of their probability masses overlap, and the solution to the classification problem is not unique. Our algorithm does not work in this case, though a recent spectral technique of Vempala and Wang [25] does apply. (However, it doesn't apply to non-spherical gaussians.)

The second problem concerns general gaussians whose probability masses do not overlap much but which appear to coalesce under random projection. For example, consider a pair of concentric gaussians that have the same axis orientation. (Of course, these axes are unknown and are not the same as the coordinate axes.) In $n - 2$ axis directions their variance is σ^2 , and in other remaining two directions their variances are $1, \sigma$ and $\sigma, 1$ respectively. If $\sigma = \Omega(\log n)$ the difference in the last two coordinates is enough to differentiate their samples with probability $1 - 1/\text{poly}(n)$. But after projection to a $O(\log n)$ dimensional subspace, this difference disappears. Hence neither distance-based clustering nor projection-based clustering seems able to distinguish their samples.

The third open problem concerns max-likelihood estimation, which seems to involve combinatorial optimization with very bizarre objective criteria once we allow non-spherical gaussians.

We suspect all the above open problems may prove difficult.

We note that Dasgupta (personal communication) has also suggested a variant of the classification problem in which the sample comes from a "noisy" gaussian. Roughly speaking, the samples come from a mixture of sources, where each source is within distance ϵ of a gaussian. We can solve this problem in some cases for small values of ϵ , but that will be subject of another paper. Broadly speaking, the problem is still open.

Acknowledgments

We thank Sanjoy Dasgupta for many helpful discussions, including drawing our attention to [21]. We also thank Laci Lovász and David McAllester for useful suggestions. We thank the anonymous referee for a quick, thorough critical reading and many important suggestions which have helped improve our presentation.

References

- [1] D. Applegate and R. Kannan. Sampling and integration of near log-concave functions. *Proceedings of the 23rd ACM Symposium on Theory of Computing*,

- ACM Press, pp. 156-163, 1991.
- [2] S. Arora and R. Kannan. Learning mixtures of arbitrary gaussians. *Proc. ACM Symposium on Theory of Computing*, ACM Press, pp. 247–257, 2001.
 - [3] J. Bourgain. Random points in isotropic convex sets. *Convex geometric analysis (Berkeley, CA, 1996)*, 53–58, Math. Sci. Res. Inst. Publ., 34, Cambridge Univ. Press, Cambridge, 1999.
 - [4] M. Charikar, S. Guha, E. Tardos, and D. Shmoys. A constant-factor approximation algorithm for the k-median problem. *Proc. 31st ACM Symposium in Theory of Computing*, 1999.
 - [5] S. DasGupta. Learning mixtures of gaussians. *Proc. IEEE Foundations of Computer Science*, 1999.
 - [6] S. Dasgupta and L. Schulman. A two-round variant of EM for gaussian mixtures. *Uncertainty in AI*, 2000.
 - [7] A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *J. Royal Statistics Soc. Ser. B*, **39**:1-38, 1977.
 - [8] M. E. Dyer, A. Frieze, R. Kannan, “A random polynomial time algorithm for approximating the volume of convex bodies,” *Journal of the ACM* **38**, pp. 1-17, 1991.
 - [9] P. Drineas, A. Frieze, R. Kannan, V. Vinay and S. Vempala, “Clustering in large graphs and matrices” *Proceedings of the ACM-SIAM Symposium on Discrete Algorithms, 1999* pp 291-299.
 - [10] Y. Freund and Y. Mansour. Estimating a mixture of two product distributions. *ACM Conference on Computational Learning Theory*, 1999.
 - [11] W. J. Hoeffding. Probability inequalities for sums of bounded random variables. *J. American Statistical Assoc*, **58**(301):13-30, March 1963.
 - [12] P. J. Huber. Projection pursuit. *Annals of Statistics*, **13**(2):435-475, June 1985.
 - [13] W. B. Johnson and J. Lindenstrauss. Extensions of Lipschitz mapping into Hilbert space. *Contemp. Math.* **26**:189-206, 1984.
 - [14] R. Kannan, G. Li, *Sampling according to the multivariate normal density*, in the Proceedings of the 37 th Annual Symposium on the Foundations of Computer Science, IEEE (1996) pp 204-213.
 - [15] B. Lindsay. *Mixture models: theory, geometry, and applications*. American Statistical Association, Virginia 1995.
 - [16] L. Lovász and M. Simonovits Mixing rate of Markov chains, an isoperimetric inequality, and computing the volume, *Proc. 31st Symposium on Foundations of Computer Science*, pp 346–355 (1990).

- [17] L. Lovász and M. Simonovits, Random walks in a convex body and an improved volume algorithm *Random Structures and Alg.* **4**, pp 359–412 1993.
- [18] N. Megiddo. On the complexity of some geometric problems in unbounded dimension, *J. Symbolic Computation* **10**:327–334, 1990.
- [19] A. Prékopa. Logarithmic concave measures with applications to stochastic programming. *Acta Sci. Math. Szeged* **32**:301–316 (1971).
- [20] A. Prékopa. On logarithmic concave measures and functions. *Acta Sci. Math. Szeged* **34**:335–343 (1973).
- [21] R. A. Redner, H. F. Walker. Mixture densities, maximum likelihood and the EM Algorithm. *SIAM Review*, **26**(2):195–239, 1984.
- [22] M. Rudelson. Random vectors in the isotropic position. *J. Func. Anal.* **164**:60–72, 1999.
- [23] D. M. Titterington, A. F. M. Smith, and U. E. Makov. *Statistical analysis of finite mixture distributions*, Wiley, 1985.
- [24] V. N. Vapnik and A. Ya Chervonenkis, *On the uniform convergence of relative frequencies of events to their probabilities*, Theory of Probability and its Applications, **16**:264–280, 1971.
- [25] S. Vempala and G. Wang, *A spectral Algorithm for Learning Mixture Models* in the Proceedings of the 43 rd Annual IEEE Symposium on the Foundations of Computer Science, 2002.